

HEALTH, EDUCATION & ATTENDANCE FOR LIFE



H.E.A.L.

FLYER FOR PARENTS



When students miss too many days of school, they fall behind and struggle to keep up with their classmates. Whether the day missed is due to illness, truancy or any other reason, the end result for the student is the same - learning time is lost. Children and adolescents will get sick at times and may need to stay at home, but we want to work with you to help minimize the number of days your student misses school.

Missed Days Add Up Quickly!

- Just a few missed days a month adds up to several school weeks missed in a year.
- Both excused and unexcused absences can make it more difficult for your child to keep up with other students, especially in math and reading.
- Kindergarten and first grade are critical for your child. Missing school during these early years makes it more difficult for children to learn in later years and they often have trouble reading by the end of third grade.

Work with Your Child and Your School

- As the parent, be strong with your child and don't let your child stay home when it is not necessary. This will help your child succeed.
- If your child has a chronic disease, make sure that the school nurse and other school staff are aware of the
 disease so they can assist your child if he or she becomes ill. Information about your child's chronic disease
 should be noted on the school emergency medical authorization form.
- For students with asthma; if your child has asthma, the school needs an Asthma Action Plan completed by his or her doctor that includes permission to use an inhaler at school. Make sure that all supplies (inhaler, spacer, etc.) needed to manage your child's asthma are at the school.
- Keep an open line of communication with school staff and teachers. The more the school knows about your child's health, the better prepared everyone will be to work together for your child.

Helpful Ideas:

- Make appointments with the doctor or dentist in the late afternoon so your child misses as little school as possible.
- If your child must miss school, make sure you get his or her homework assignments and follow up to see that the work is completed and turned in.
- Call the school as soon as you know your child will be absent and tell school staff why your child will be out and for how long.
- Be prepared to get a doctor's note when requested by school personnel.
- If you need medical advice after business hours, most doctors' offices have answering services 24 hours a
 day to assist you.
- If your child has an emergency, call 911.

For additional information contact Your school nurse!



WHEN SHOULD I SEND MY CHILD TO SCHOOL?

The suggestions below are for children 5 to 18 years of age. Recommendations may be different for infants and younger children.

| Symptoms and Illnesses | rent for infants and younger children. Should My Child Go To School? |
|--|---|
| | Should My Child Go to School? |
| Parent or other family member is sick, stressed, hospitalized | YES - If you are sick, your child still needs to attend school. Your illness does not excuse your child from attending. We are all sick at times so plan ahead for these days. Get a neighbor, relative, or spouse to take your child to school and pick him or her up. |
| Chronic Diseases (Asthma, Diabetes, Sickle Cell, Epilepsy, etc.) Chronic disease is a long lasting condition that can be controlled but not cured. | YES - Your child should attend school. School personnel are trained to assist your child with his or her chronic disease and associated needs. Talk to the school nurse. |
| Child Doesn't Want to go to School | YES - You should keep your child in school, but try to determine |
| Frequent crying, fear, anger, not wanting to socialize, behavior change, stomach ache, nausea (These can be signs of depression, anxiety, or fear) | what is causing the changes. Talk to school personnel and consult a health care provider. Your child may be experiencing bullying or trauma, may be behind in his or her school work or not getting along with others. These and other issues may require your or the school's attention. |
| Cold Symptoms Stuffy nose/runny nose, sneezing, mild cough | YES - If your child is able to participate in school activities send him or her to school. Most children with cold symptoms attend school. |
| Conjunctivitis (Pink Eye) The white of the eye is pink | YES - Your child can attend school, but call a health care provider to prescribe medication/treatment if it does not go away in 4 or 5 days. |
| Head Lice | YES - Your child can be in school if he or she has had an initial |
| Intense itching of the head; may feel like something is moving | treatment of shampooing of hair with a product for lice. |
| Strains, Sprains, and Pains | YES - If there is no known injury and your child is able to function (walk, talk, eat) he or she should be in school. If pain is severe or doesn't stop, consult a health care provider. |
| Menstrual Issues | YES - Most of the time menstrual (periods) issues should not be a problem. If they are severe and interfering with your daughter attending school, consult with a health care provider. |
| Fever Fever usually means illness, especially if your child has a fever of 100 or higher as well as other symptoms like behavior change, rash, sore throat, vomiting etc. | NO - If your child has a fever of 100 or higher keep them at home until his or her fever is below 100 for 24 hours without the use of fever reducing medication. If the fever does not go away after 2-3 days or is 102 or higher, you should consult a health care provider. |
| Diarrhea Frequent, loose or watery stool may mean illness but can also be caused by food and medication | NO - If, in addition to diarrhea, your child acts ill, has a fever or is vomiting, keep him or her at home. If stool is bloody, if the child has abdominal pain, fever or vomiting, you should consult a health care provider. |
| Vomiting | NO - Keep your child at home until the vomiting has stopped 24 hours. If vomiting continues, contact a health care provider. |
| Child has vomited 2 or more times in a 24 hour period | NO Variance bild have and a first fit |
| Coughing Severe, uncontrolled, rapid coughing, wheezing, or difficulty breathing | NO - Keep your child home and contact a health care provider. Asthma - if symptoms are due to asthma, provide treatment according to your child's Asthma Action Plan and when symptoms are controlled send your child to school. |
| Rash With Fever | NO - If a rash spreads quickly, is not healing, or has open weeping wounds, you should keep your child at home and have him or her seen by a health care provider. Keep child at home for 24 hours after an antibiotic is begun. |
| Sore Throat | NO - Keep your child at home for the first 24 hours after an antibiotic is begun. |
| With fever, stomach ache, and red, swollen tonsils | , |

This information is based upon recommended guidelines from reliable sources including the Centers for Disease Control (CDC), American Academy of Pediatrics, and Public Health Association. It has been adapted with permission from the Alameda County California District Attorney's Office.